

Passing a bottleneck, the fish – bubble analogy

Encadrement : Aurélie Dupont & Benjamin Dollet Laboratoire Interdisciplinaire de physique, Grenoble

> <u>Benjamin.dollet@univ-grenoble-alpes.fr</u> Aurelie.dupont@univ-grenoble-alpes.fr

Collective movements are a fascinating phenomenon that can be observed in living organisms at different scales, from bacteria to human crowds. We are interested in the case of schools of fish and more particularly in their evacuation through an opening of a size comparable to the size of the fish. We have recently shown that the escape of fish is analogous to that of bubbles[1]. Indeed, the fish outflow (fish/s) is well fitted by the Beverloo law used for bubbles in a similar situation by Bertho et al.[2]. We have modeled the fish as surrounded by a cognitive bubble that governs their outflow statistics (Figure, left).

We would like to further explore this analogy and its limits. To this end, we have recently built an experiment (Figure, right) allowing the evacuation of bubbles through one or two slits. We can vary the size of the opening, the evacuation stress and the size of the bubbles. The aim of the internship will be to analyze the statistics of evacuation time intervals, in the same way as we did for fish, and see whether the model proposed for fish is valid for bubbles.

Fish current L_c

(Left) Model of fish surrounded by a cognitive bubble while passing through a bottleneck. (Right) Preliminary results of bubbles evacuating through an opening.

The physical questions are many and open: how intermittent is bubble evacuation? Can we

observe blocking phenomena such as those seen with active particles or grains? What are the interactions between slits and their consequences on bubble evacuation rates? Do all the bubbles flow like grains in an hourglass, liquid in a clepsydra, or something else entirely?

Based on preliminary work, the trainee will have to finalize the experimental protocol by identifying the most controlled bubble injection conditions possible, and carry out the various experiments with the bubbles. He/she will then analyze the images obtained and take part in the modeling.

The internship can be extended into a PhD.

- [1] Larrieu, R., Moreau, P., Graff, C., Peyla, P., & Dupont, A. (2023). Fish evacuate smoothly respecting a social bubble. Scientific Reports 13 (1), 10414
- [2] Bertho, Y., Becco, C., & Vandewalle, N. (2006). Dense bubble flow in a silo: An unusual flow of a dispersed medium. *Physical Review E*, 73(5), 056309.





