

## New and Notable

### Pack it up, Pack it in: Unraveling H-NS Mediated Genome Packaging

Gordon S. Freeman<sup>†</sup>  
and Juan J. de Pablo<sup>‡\*</sup>

<sup>†</sup>Department of Chemical and Biological  
Engineering, University of Wisconsin-  
Madison, Madison, Wisconsin; and

<sup>‡</sup>Institute for Molecular Engineering,  
University of Chicago, Chicago, Illinois

How are bacterial genomes compacted? Nucleoid-associated proteins have been discovered that bind to the circular genome in either a sequence-dependent or sequence-independent manner, thereby resulting in a compact nucleoid that can be confined within the bacteria and be accessed by the cellular machinery required to transcribe the genome (1,2). Dame and co-workers performed a number of key experiments that elucidated the binding behavior of a particular nucleoid-associated protein, histone-like nucleoid structuring protein (H-NS), and its role in confining the genome. Their work provided a model for nucleoid condensation in which H-NS protein binds randomly to one strand and, upon meeting another strand of DNA, induces condensation through *trans* binding (3). Through optical-tweezer experiments, they were able to further characterize the kinetics and thermodynamics of this process (4). However, open questions remained as the work demonstrating condensation of DNA by H-NS was performed in two dimensions on a mica surface and neither the effect of this spatial confinement nor the nature of the DNA-surface interaction itself were fully addressed.

Seeking to better understand the experimental findings of Dame et al., in their article Joyeux and Vreede (5)

develop a representation of DNA and H-NS dimers consisting of bead-spring constructs. Through a careful choice of parameters, they closely reproduce the experimentally observed binding behavior of H-NS to DNA. They are able to then use this simple model to help elucidate the rich physics observed in experiments probing H-NS mediated genome condensation. In particular, their work gives insight into the nature of the H-NS dimer itself and its effect on genome condensation. In the work of Wiggins et al. (6), protein-mediated bridging is discussed in the context of two possible structural motifs, an H-NS linker domain (residues 65–89) that is either rigid or flexible. In the language of Joyeux and Vreede, this corresponds to an H-NS with a large or small value of  $G$  (the H-NS dimer bending rigidity), respectively. Coarse-grained simulations enable Joyeux and Vreede to test a hypothesis such as that of Wiggins et al. by simply assigning the dimer different physical parameters and observing the resulting change in behavior. They are able to demonstrate that a difference of ~20% in binding affinity in the *cis* configuration (the result of a factor of two change in the H-NS bending rigidity) is sufficient to fundamentally change the dynamics of the condensed nucleoid, with the more flexible case resulting in a fluctuating, dynamic structure exhibiting open loops (more *cis* binding) whereas the less flexible case results in a more compact globular structure that changes little over time (increased *trans* binding). Joyeux and Vreede do not propose a precise value for the H-NS bending rigidity, as the experimental data to which they compare are difficult to interpret (primarily nucleoids condensed on a two-dimensional mica surface (3)); their work, however, is of fundamental significance in that it points the way toward experiments that might better elucidate the true nature of H-NS and its role in nucleoid condensation. Indeed the marked difference in the

radius of gyration of three-dimensional condensed nucleoids reported in their work, driven by differences in *cis* binding affinity, represents a quantity that can be directly accessed in experiment to make progress in the problem of H-NS mediated genome condensation and the role played by H-NS flexibility and binding affinity therein.

Looking forward in terms of developing molecular-level models to further understand protein-mediated genome compaction, the growing availability of advanced sampling techniques and reliable models provides the biophysics community with tools at every level of molecular description. The coarse-grained representation adopted by Joyeux and Vreede has provided valuable insights. Building on their results, future studies should aim to address potentially interesting details of the interaction between H-NS and DNA. Experimental data indicate that this interaction is not, in fact, entirely sequence-agnostic (as the Joyeux and Vreede representation assumes), but instead exhibits a preference for AT-rich genomic regions (7). Specifically, the minor groove width in these AT-rich regions is thought to be optimal for H-NS binding. Coarse-grain DNA (8,9) and protein (10) models have been developed that are capable of exploring such shape-dependent protein-nucleic acid interactions; approaching the problem at this scale may yield rich information regarding H-NS/DNA interactions. A particular detail that remains to be addressed is the effect of the ionic environment on the interaction of H-NS with the genome. In earlier work, Vreede and Dame used molecular simulations to demonstrate that the conformation of the dimerization domain of H-NS may indeed be sensitive to ionic conditions, with the parallel dimer increasing in stability with increasing salt (11). Experimental

Submitted February 11, 2013, and accepted for  
publication February 25, 2013.

\*Correspondence: [depablo@uchicago.edu](mailto:depablo@uchicago.edu)

Editor: Kathleen Hall.

© 2013 by the Biophysical Society  
0006-3495/13/04/1397/2 \$2.00



evidence also indicates that altering the ionic environment can fundamentally change the nature of the resulting H-NS/DNA complexes and promote either the *cis* or *trans* binding regime (12,13). Coarse-grained molecular models are uniquely suited to explore this wide range of environmental conditions and assess their impact on the mechanism of H-NS mediated nucleoid condensation.

A particularly exciting idea that could now be addressed is that AT-rich sequences that promote H-NS bridging behavior act as domain barriers in bacterial genomes (1,14). Interestingly, the acquisition of genomic islands, generally AT-rich, is believed to potentially disrupt the three-dimensional structure of the nucleoid through the recruitment of H-NS proteins and decrease the fitness of the bacterial cell (14). This notion that stochastic addition of H-NS binding domains may drastically alter the organization of the genome could be directly probed in coarse-grained simulations in which sequence-specificity in H-NS binding is taken into account.

In their work, Joyeux and Vreede provide a compelling example of how

a simple and elegant molecular model can be used to provide profound insight into an otherwise complex biophysical problem. In addition to highlighting the large impact that small changes in DNA-protein interactions can have, their work suggests key experiments that, in due course, will provide a clearer picture of H-NS mediated genome condensation.

## REFERENCES

1. Wang, W., G. W. Li, ..., X. Zhuang. 2011. Chromosome organization by a nucleoid-associated protein in live bacteria. *Science*. 333:1445–1449.
2. Luijsterburg, M. S., M. F. White, ..., R. T. Dame. 2008. The major architects of chromatin: architectural proteins in bacteria, archaea and eukaryotes. *Crit. Rev. Biochem. Mol. Biol.* 43:393–418.
3. Dame, R. T., C. Wyman, and N. Goosen. 2000. H-NS mediated compaction of DNA visualized by atomic force microscopy. *Nucleic Acids Res.* 28:3504–3510.
4. Dame, R. T., M. C. Noom, and G. J. L. Wuite. 2006. Bacterial chromatin organization by H-NS protein unraveled using dual DNA manipulation. *Nature*. 444:387–390.
5. Joyeux, M., and J. Vreede. 2013. A model of H-NS mediated compaction of bacterial DNA. *Biophys. J.* 104:1615–1622.
6. Wiggins, P. A., R. T. Dame, ..., G. J. Wuite. 2009. Protein-mediated molecular bridging: a key mechanism in biopolymer organization. *Biophys. J.* 97:1997–2003.
7. Gordon, B. R. G., Y. Li, ..., J. Liu. 2011. Structural basis for recognition of AT-rich DNA by unrelated xenogeneic silencing proteins. *Proc. Natl. Acad. Sci. USA*. 108:10690–10695.
8. Sambriski, E. J., D. C. Schwartz, and J. J. de Pablo. 2009. A mesoscale model of DNA and its renaturation. *Biophys. J.* 96:1675–1690.
9. Ortiz, V., and J. J. de Pablo. 2011. Molecular origins of DNA flexibility: sequence effects on conformational and mechanical properties. *Phys. Rev. Lett.* 106:238107.
10. Takada, S. 2012. Coarse-grained molecular simulations of large biomolecules. *Curr. Opin. Struct. Biol.* 22:130–137.
11. Vreede, J., and R. T. Dame. 2012. Predicting the effect of ions on the conformation of the H-NS dimerization domain. *Biophys. J.* 103:89–98.
12. Amit, R., A. B. Oppenheim, and J. Stavans. 2003. Increased bending rigidity of single DNA molecules by H-NS, a temperature and osmolarity sensor. *Biophys. J.* 84:2467–2473.
13. Liu, Y., H. Chen, ..., J. Yan. 2010. A divalent switch drives H-NS/DNA-binding conformations between stiffening and bridging modes. *Genes Dev.* 24:339–344.
14. Noom, M. C., W. W. Navarre, ..., R. T. Dame. 2007. H-NS promotes looped domain formation in the bacterial chromosome. *Curr. Biol.* 17:R913–R914.