

Nonlocal stationary probability distributions and escape rates for an active Ornstein–Uhlenbeck particle

Eric Woillez¹, Yariv Kafri¹ and Vivien Lecomte²

¹ Department of Physics, Technion, Haifa 32000, Israel

² Université Grenoble Alpes, CNRS, LIPhy, F-38000 Grenoble, France

E-mail: eric.woillez@neel.cnrs.fr

Received 10 December 2019

Accepted for publication 28 February 2020

Published 9 June 2020



Online at stacks.iop.org/JSTAT/2020/063204

<https://doi.org/10.1088/1742-5468/ab7e2e>

Abstract. We evaluate the steady-state distribution and escape rate for an active Ornstein–Uhlenbeck particle (AOUP) using methods from the theory of large deviations. The calculation is carried out both for small and large memory times of the active force in one-dimension. We compare our results to those obtained in the literature about colored noise processes, and we emphasize their relevance for the field of active matter. In particular, we stress that contrary to equilibrium particles, the invariant measure of such an active particle is a non-local function of the potential. This fact has many interesting consequences, which we illustrate through two phenomena. First, active particles in the presence of an asymmetric barrier tend to accumulate on one side of the potential—a ratchet effect that was missing in previous treatments. Second, an active particle can escape over a deep metastable state without spending any time at its bottom.

Keywords: active matter, large deviations in non-equilibrium systems, boundary layers

Supplementary material for this article is available [online](#)

Contents

1. Introduction	2
2. The approach and some implications of the results	6
2.1. Summary of the large deviation framework.....	6
2.2. The small τ limit	8
2.2.1. Instanton path for the harmonic potential	9
2.2.2. A ratchet	10
2.3. The large τ limit	11
2.3.1. Hopping over metastable states	14
3. AOUP in the small τ limit	15
3.1. Expansion order by order.....	15
3.1.1. Zeroth order	16
3.1.2. First order	17
3.1.3. Second order	18
4. AOUP in the large τ limit	20
4.1.1. Bulk solution.....	21
4.1.2. Boundary layer solution	23
4.2. The quasipotential for $x < x_{cr}$	29
4.2.1. Bulk solution.....	29
4.2.2. Boundary-layer solution	29
5. Perspectives	31
Acknowledgments	31
Appendix. Mean escape time from a metastable state	32
References	33

1. Introduction

The active Ornstein–Uhlenbeck process (AOUP in the following) is defined in one dimension through the equations of motion

$$\begin{cases} \dot{x} = v - U'(x), \\ \tau \dot{v} = -v + \sqrt{2D}\xi(t). \end{cases} \quad (1)$$

The model can be interpreted as describing the overdamped dynamics of a particle with position x in an external potential U (with the mobility set to one) and driven

by a Gaussian noisy velocity v with zero average and a time–time correlation function $\langle v(t)v(t') \rangle = \frac{D}{\tau} e^{-|t-t'|/\tau}$. First studied in the context of Langevin equations with memory [1–10], it has received much attention recently when it became one of the canonical models used to describe the motion of active particles [11–16]. Active particles are a particular class of systems where each particle consumes energy from its environment in order to self-propel. This AOUP model captures this through the stochastic active force v which has a persistence time τ that breaks time reversal symmetry, and therefore drives the system out of equilibrium: when τ goes to zero, one recovers the model of equilibrium Brownian particles, but as long as $\tau > 0$, there is a nonzero entropy production (of order τ^2 as $\tau \rightarrow 0$ [15]).

The model can be solved exactly when U is a harmonic potential. Then, it can be shown that the AOUP process can be mapped to an equilibrium process. The invariant measure in the potential and the escape rate, assuming that the particle escapes when it reaches a given potential height, can be computed exactly (see for example, [17]). For general potentials most of the work has focused on the limit $\tau \rightarrow 0$. There, approximation techniques, specifically the unified colored noise approximation (UCNA) and Fox approximation, have been employed in order to obtain the steady-state distribution [9, 10]. These show that to first order in τ , the AOUP is similar to an equilibrium system but with an effective *local* potential whose form can be obtained and which depends on the second spatial derivative of U [13]. The term *local* emphasizes the fact that the steady-state distribution is a local function of the potential and its derivatives. The result, however, is restricted to convex potentials, with the additional constraint that the ratio $\sqrt{\langle v^2 \rangle} = \sqrt{D/\tau}$ remains large compared to typical values of the force U' . To this end, bistable potentials which cannot be treated using the UCNA and Fox approximations, have been studied numerically in [16] where an alternative approximation scheme in the small τ limit was developed.

More recently the escape problem of an AOUP from a general trap was considered in the large τ limit [17]. In this regime, it was shown that the behavior changes in a drastic manner from the leading order in τ result at small τ . Specifically, the escape rate from the trap does not depend on the maximum difference in the value of an effective potential, but on the *maximal force* exerted on the particle inside the trap. This suggests that the steady-state distribution takes on a very different form when orders higher than τ are accounted for. The following paper pursues this line of thought by calculating both the steady-state distribution and the closely related problem of the escape rate from a trap, in the small D limit, both to order τ^2 and in the limit of $\tau \rightarrow \infty$. We show that in both limits the physics becomes completely different from that revealed by the leading order in τ result. In particular, the steady-state distribution is now a *non-local* function of the potential U . This is crucial since, as we illustrate, fundamental non-equilibrium physical processes such as ratchet effects are only displayed when the non-locality of the steady state appears. These, for example in the seminal work of [18–20], are considered one of the hallmarks of active matter.

To obtain these results, as stated above and in contrast to previous studies of the AOUP model, we focus on the small-noise limit. Namely, we consider the problem when D , the amplitude of the noise which drives the variations in v , is small. The use of a small noise asymptotic analysis to study the problem has been first applied about three

decades ago in the context of Brownian motion subject to colored noise³. In particular, a path integral formulation was derived, despite the non-Markovian evolution of the position [1, 2, 4–6, 21]. These formulations in turn allow one to study the $D \rightarrow 0$ asymptotic regime through an optimal path approach. In particular, they allow for the computation of escape rates [4, 7, 21] including the sub-dominant prefactor [8] and even accounting for inertia [22] (see also [2] for solutions of specific models or [23] for other correlated noise). We also note an early work by Kłosek-Dygas *et al* [3] who, by means of a Fokker–Planck approach (and in a small- τ expansion), obtained many of the results derived later through a path-integral approach. Below we compare in detail our results to those obtained previously emphasizing technical differences, places where we go beyond the early results, and putting them in the context of active matter. As we show the focus on phenomena related to active matter allows us to uncover new phenomena which were not discussed in the early works.

To study the $D \rightarrow 0$ asymptotics we apply recent techniques from the theory of large deviations [24]. These enable us to relate the escape problem and the steady-state distribution problem using standard results. In particular, if we consider a metastable state x_0 and its basin of attraction \mathcal{B}_0 both the mean escape time $\langle T_{\text{esc}} \rangle$ and the invariant measure ρ within \mathcal{B}_0 can be asymptotically computed from a single function Φ_τ called the quasipotential as follows

$$\begin{cases} \frac{\rho(x)}{\rho(x_0)} & \underset{D \rightarrow 0}{\asymp} e^{-\frac{\Phi_\tau(x)}{D}}, \\ \langle T_{\text{esc}} \rangle & \underset{D \rightarrow 0}{\asymp} e^{\frac{\Phi_\tau(x_s)}{D}}. \end{cases}$$

Here x_s is the saddle point that needs to be crossed for the particle to escape from the basin of attraction of x_0 . The notation $f \underset{D \rightarrow 0}{\asymp} g$ means logarithmic equivalence between the functions f and g , that is $\log f \underset{D \rightarrow 0}{\sim} \log g$. Using standard tools we compute the quasipotential Φ_τ in both limits $\tau \rightarrow 0$ (to order τ^2) and $\tau \rightarrow +\infty$, without any restriction on U . Our main findings are:

- (a) In the limit $\tau \rightarrow 0$, we find the asymptotic expansion

$$\Phi_\tau(x) \underset{\tau \rightarrow 0}{\sim} U(x) - U(x_0) + \tau \frac{|U'(x)|^2}{2} - \frac{\tau^2}{2} \int_{x_0}^x |U'(y)|^2 U'''(y) dy + O(\tau^3). \quad (2)$$

The correction of order τ^2 in Φ_τ is the first **nonlocal term** of the invariant measure. We interpret this contribution in terms of long-range interactions in the system. These, as we show, lead to a ratchet effect where particles are driven in the presence of an asymmetric potential (see section 2.2.2 and figure 2) [20]. We note that non-local interactions have been already observed in active fluids [25] and therefore seem to be generic for this class of systems. The non-local contribution, interestingly, is absent for a harmonic potential. Our method can further be used to derive an

³In an earlier version of the paper we were not aware of these pioneering works. We thank an anonymous referee for bringing our attention to references [4, 6, 7] and apologize to their authors.

expansion for Φ_τ to any order in τ . We note from a technical point of view that the expansion (2) in powers of τ is non-intuitive: because of the existence of a boundary layer in the instanton calculation used to evaluate $\Phi_\tau(x)$, a standard perturbative procedure fails and the calculation has to be carried out with some care. The singular behavior of the instanton also brings an insight on how an active optimal trajectory differs from an equilibrium one.

Comparing to earlier works, the $O(\tau)$ contribution to (2) was derived by Wio *et al* in [21] using a path integral approach. The escape rate implied by (2) (i.e. for $x = x_s$) was derived in [4, 7] at order τ^2 and higher using path integrals. However, in the latter case the optimal path is not singular, and the perturbation is organized in even powers of τ . The methods of [4, 7] therefore allows one to determine the escape rate but not the steady-state quasipotential for any x . However, as discussed at the end of section 3, we note that expression (2) can be deduced from a result of referene [3] which uses a Fokker–Planck approach. This provides an alternative way to derive (2), but one does not uncover the singular nature of the instanton. The approach we follow gives an insight on the typical trajectory taken by the active particle to reach an arbitrary point.

- (b) The limit $\tau \rightarrow +\infty$ displays a different behavior. The quasipotential should be seen as a functional of the force field $F(x) = -U'(x)$, rather than a functional of the potential. We show that the quasipotential has a singularity at the inflection point x_{cr} where $U''(x_{\text{cr}}) = 0$. Before the inflection point, Φ_τ is a local function of the force, and is given by

$$\text{for } x < x_{\text{cr}}, \quad \Phi_\tau(x) \underset{\tau \rightarrow +\infty}{\sim} \tau \frac{|U'(x)|^2}{2} + \frac{|U'(x)|^2}{2U''(x)} + O(1/\tau). \quad (3)$$

However, beyond the inflection point, it is entirely controlled by the vicinity of x_{cr}

$$\text{for } x > x_{\text{cr}}, \quad \Phi_\tau(x) \underset{\tau \rightarrow +\infty}{\sim} \tau \frac{|U'(x_{\text{cr}})|^2}{2} + \tau^{1/3} \frac{C|U'(x_{\text{cr}})|^2}{\left(\sqrt{\frac{1}{2}|U'''(x_{\text{cr}})|}|U'(x_{\text{cr}})|\right)^{2/3}} + O(\tau^0), \quad (4)$$

where $C \approx 0.8120$ is a constant that can be computed numerically from the solution of a non-dimensional equation (see equations (74) and (75) and section 4.1.2). This demonstrates again the non-locality of the problem. The result implies that in the large τ limit, the potential barrier seen by the active particle is controlled by forces. For example, in the escape problem the particle has to overcome the maximal force $F_{\text{max}} = U'(x_{\text{cr}})$, as opposed to the equilibrium problem where maximal potential is the bottleneck. The proof of equation (4) further reveals that the dynamical system (1) has a bifurcating slow manifold that is used by the active particle to switch between the metastable states. This particular kind of transition paths has been already observed numerically for some multiscale systems in [26]. This gives rise to interesting new phenomena, which we discuss, such as hopping over metastable states (see especially section 2.3.1 and the supplementary movies, available online at ([stacks.iop.org/JPhysJSTAT/2020/063204/mmedia](https://doi.org/10.1088/1742-5468/ab7e2e))).

In comparison to previous works we note that the $\tau^{1/3}$ scaling of the correction (4) was hinted at in reference [4]. Expression (4), in the case $x = x_{\text{cr}}$ useful in the escape problem, was derived in [7] through a path-integral approach, but again the singular nature of the instanton (that appears only for $x \neq x_{\text{cr}}$) was not identified. Finally, we comment that the steady-state quasipotential cannot be inferred from the approach of reference [7].

The paper is organized as follows: for completeness we first give in section 2 a summary of the approach used for the calculations, before discussing their consequences. The detailed derivations of the results are given in sections 3 and 4.

2. The approach and some implications of the results

2.1. Summary of the large deviation framework

In this section we describe the approach used to evaluate the quasipotential Φ_τ . For completeness, the relation between the mean escape time and the quasipotential is recalled in the appendix. In what follows, we assume for simplicity that $\rho(x_0) = 1$, the generalization being straightforward. First, note that the invariant measure can be expressed in terms of the transition probability of the AOUP as

$$\rho(x) = \lim_{T \rightarrow -\infty} \int_{-\infty}^{+\infty} P(x, v, 0 | x_0, 0, -T) dv,$$

where $P(x, v, 0 | x_0, 0, -T)$ is the probability to be at (x, v) at time $t = 0$ starting initially at the stationary point $(x_0, 0)$ at $t = -T$. In this paper, we use both the Lagrangian and the Hamiltonian formalism, and we briefly review both in the following.

Lagrangian formalism: the transition probability can then be expressed as an Onsager–Machlup path integral [27]. This gives for the invariant measure

$$\rho(x) = \int \mathcal{D}[x(t), v(t)] e^{-\frac{1}{\tau} \mathcal{A}[x, v]}, \quad (5)$$

with

$$\mathcal{A}[x, v] = \int_{-\infty}^0 \mathcal{L}(x, v, \dot{x}, \dot{v}) dt \quad (6)$$

$$\mathcal{L}(x, v, \dot{x}, \dot{v}) = \begin{cases} \frac{1}{4}(\tau \dot{v} + v)^2 & \text{if } v = \dot{x} + U'(x) \\ +\infty & \text{otherwise.} \end{cases} \quad (7)$$

Using the constraint on v the Lagrangian can be equivalently written as

$$\mathcal{L}(x, \dot{x}, \ddot{x}) = \frac{1}{4}(\dot{x} + U'(x) + \tau(\ddot{x} + \dot{x}U''(x)))^2. \quad (8)$$

The instanton trajectory is found by solving the Euler–Lagrange equation with the boundary conditions

$$\begin{aligned} x(t) &\xrightarrow{t \rightarrow -\infty} x_0; \quad v(t) \xrightarrow{t \rightarrow -\infty} 0 \\ x(0) &= x; \quad \dot{x}(0) = 0. \end{aligned} \quad (9)$$

The first line implies that the particle starts at the stationary state, while the second assures that it arrives at x at time zero without overshooting. So far, expression (5) is formally exact (with a proper interpretation/discretization of the action, see e.g. [28]). In the limit $D \rightarrow 0$, one can use a saddle-point approximation to get

$$\int \mathcal{D}[x(t), v(t)] e^{-\frac{1}{D} \mathcal{A}[x, v]} \underset{D \rightarrow 0}{\simeq} e^{-\frac{1}{D} \Phi_\tau(x)},$$

where $\Phi_\tau(x)$ is the quasipotential.

Hamiltonian formalism: the easiest path for obtaining the Hamiltonian formalism is through the Martin–Siggia–Rose–Janssen–De Dominicis approach [29–33]. To do so we rewrite the probability by reexpressing delta functions as

$$\rho(x) = \left\langle \int \mathcal{D}[x(t), v(t), P_v(t), P_x(t)] e^{-\frac{1}{D} \int_{-\infty}^0 \{P_x(\dot{x} - v + U') + P_v(\tau \dot{v} - v - \sqrt{2D} \xi(t))\} dt} \right\rangle_\xi, \quad (10)$$

which after averaging over the noise ξ gives

$$\rho(x) = \int \mathcal{D}[x(t), v(t), P_v(t), P_x(t)] e^{-\frac{1}{D} \int_{-\infty}^0 \{P_x \dot{x} + P_v \dot{v} - H[P_x, P_v, x, v]\} dt}, \quad (11)$$

where the Hamiltonian is given by

$$H(x, v, P_x, P_v) = (v - U'(x)) P_x - \frac{v P_v}{\tau} + \frac{P_v^2}{\tau^2}. \quad (12)$$

The instanton is then given by a solution of Hamilton’s equations

$$\begin{cases} \dot{x} = v - U'(x) \\ \dot{v} = -\frac{v}{\tau} + 2\frac{P_v}{\tau^2} \end{cases} \quad \text{and} \quad \begin{cases} \dot{P}_x = U''(x) P_x \\ \dot{P}_v = \frac{P_v}{\tau} - P_x \end{cases}, \quad (13)$$

with the boundary conditions

$$x(t) \xrightarrow{t \rightarrow -\infty} x_0; \quad v(t), P_x(t), P_v(t) \xrightarrow{t \rightarrow -\infty} 0. \quad (14)$$

$$P_v(0) = 0 \quad (15)$$

The boundary condition on $P_v(0)$ can be deduced by noting that $P_v = \partial_v S$, with S the action corresponding to H , and that we are minimizing the action with respect to the final velocity. It is equivalent to the Lagrangian one, equation (9). The boundary conditions at $t = -\infty$ results from demanding that we start at the stationary point.

Finally, note that the Hamiltonian has no explicit time dependence, ‘energy’ is conserved along the optimal path. Evaluating the Hamiltonian (12) at $t = -\infty$ gives the constraint

$$H(x, v, P_x, P_v) = 0. \quad (16)$$

With the above we now turn to discuss the small and large τ expansions.

2.2. The small τ limit

We now explain how the value of the quasipotential at the saddle point $\Phi_\tau(x_s)$ can, for the escape problem, be evaluated to order τ^2 using a standard perturbation theory [4, 7]. Note that this approach, as we discuss in section 3.1, in fact fails when the final point x is not a saddle point x_s , i.e. for the computation of the invariant measure $\Phi_\tau(x)$ at an arbitrary point x . We will show how the perturbation scheme has to be modified in this case. For clarity, these subtleties will be illustrated for a harmonic potential $U(x) = \frac{1}{2}kx^2$ in section 2.2.1, where $\Phi_\tau(x)$ is computed exactly. In what follows we always take $U(x_0) = 0$.

To proceed, we first expand the square in the action (6) and (7) so as to write

$$\mathcal{A}[x] = \mathcal{A}_0[x] + \tau \mathcal{A}_1[x] + \tau^2 \mathcal{A}_2[x].$$

Note that equation (7) implies that $\mathcal{A}_1[x] = \frac{1}{2} \int_{-\infty}^0 \dot{v} v dt = \frac{1}{4} v(0)^2$ is a boundary term. Furthermore, since we are looking for action minimizers which reach the saddle x_s , the fluctuation path \tilde{x} (or optimal path) has to reach x_s with zero velocity. This can be directly seen from equation (16) with the boundary condition equation (15). Therefore, for x_s a saddle, $\mathcal{A}_1[\tilde{x}] = 0$. Next, using equation (7)

$$\begin{cases} \mathcal{A}_0[x] = \frac{1}{4} \int_{-\infty}^0 v^2 dt, \\ \mathcal{A}_2[x] = \frac{1}{4} \int_{-\infty}^0 \dot{v}^2 dt. \end{cases} \quad (17)$$

As the action expansion has no terms proportional to τ , we can expand the fluctuation path $\tilde{x}(t)$ and the quasipotential $\Phi_\tau(x_s)$ in powers of τ^2

$$\tilde{x}(t) = \tilde{x}_0(t) + \tau^2 \tilde{x}_2(t) + O(\tau^4) \quad (18)$$

$$\Phi_\tau(x_s) = \Phi_0(x_s) + \tau^2 \Phi_2(x_s) + O(\tau^4) \quad (19)$$

and solve the minimization problem up to order τ^2 . The *zeroth* order is given by the equilibrium fluctuation path

$$\begin{cases} \dot{\tilde{x}}_0(t) = U'(\tilde{x}_0(t)), \\ \mathcal{A}_0[\tilde{x}_0] = U(x_s), \end{cases} \quad (20)$$

and the mean escape time is given by the classical Arrhenius law, with the quasipotential

$$\Phi_0(x_s) = U(x_s) - U(x_0).$$

The second order correction to the quasipotential is given by

$$\Phi_2(x_s) = \int \frac{\delta \mathcal{A}_0}{\delta x} [\tilde{x}_0(t)] \tilde{x}_2(t) dt + \mathcal{A}_2[\tilde{x}_0(t)].$$

Because $\tilde{x}_0(t)$ is a minimizer of \mathcal{A}_0 , the first term vanishes and we are left with

$$\Phi_2(x_s) = \mathcal{A}_2[\tilde{x}_0(t)] = \int_{-\infty}^0 (U' U'')^2 dt, \quad (21)$$

where the integral has to be evaluated along the equilibrium fluctuation path (20). Finally, using $dt = \frac{dx}{|U'|}$, see equation (20), we obtain

$$\Phi_2(x_s) = \int_{x_0}^{x_s} |U'(y)| (U''(y))^2 dy. \quad (22)$$

This relation gives the correction to order τ^2 for the Arrhenius mean escape time, as $\langle T_{\text{esc}} \rangle \underset{D \rightarrow 0}{\asymp} e^{\frac{\Phi_\tau(x_s)}{D}}$ with the expansion given in equation (19).

As mentioned above the perturbative procedure described above and the result of equation (22) (known since the works of Bray, MacKane and collaborators [4, 7]) **cannot** be used, in a straightforward manner, to compute the perturbative corrections to the invariant measure. The reason is that the fluctuation path which reaches a point x that is not a saddle (i.e. $x \neq x_s$) displays a boundary layer of size τ close to $t = 0$. This peculiarity of the fluctuation path gives local contributions to the quasipotential to any orders in τ , that only vanish at $x = x_s$. This effect is already present for the exactly solvable case of harmonic potential as illustrated in section 2.2.1. The full computation of the fluctuation path and the quasipotential for a general potential up to order τ^2 is rather lengthy, and is detailed in section 3. The final result was announced in equation (2). As emphasized previously, equation (2) shows that the second order correction to the quasipotential is *non-local*. Several hallmarks characterizing active systems, such as ratchet currents, originate from such contributions. We illustrate this on the ratchet example in section 2.2.2.

2.2.1. Instanton path for the harmonic potential. To illustrate the subtle boundary layer that emerges in the instanton path for $\tau > 0$, we consider the AOUP in a harmonic potential $U(x) = \frac{1}{2} k x^2$. In this case, both the fluctuation path and the quasipotential can be computed exactly. Interestingly, all terms to order larger than τ in the expansion vanish. Therefore, the problem reduces to an effective equilibrium Brownian particle with an ‘effective temperature’ that depends on τ and the stiffness of the trap k (see e.g. [17]).

To see this, define the variable $p = v - kx$, so that equation (1) become

$$\begin{cases} \dot{x} = p, \\ \dot{p} = -\left(\frac{1}{\tau} + k\right)p - \frac{k}{\tau}x + \sqrt{\frac{2D}{\tau^2}}\xi(t). \end{cases}$$

Then, we can identify a friction coefficient $\gamma = \frac{1}{\tau} + k$, and effective potential $U_{\text{eff}}(x) = \frac{kx^2}{2\tau}$ so that

$$\begin{cases} \dot{x} = p, \\ \dot{p} = -\gamma p - \nabla U_{\text{eff}} + \sqrt{\frac{2\gamma D}{\tau + k\tau^2}} \xi(t). \end{cases} \quad (23)$$

equation (23) are exactly the equations of an underdamped equilibrium Brownian particle in a potential $U_{\text{eff}}(x)$ and a temperature $T_{\text{eff}} = \frac{D}{\tau + k\tau^2}$. The problem can therefore be solved exactly:

- (a) The **mean escape time** satisfies the Arrhenius law

$$\langle T_{\text{esc}} \rangle \asymp e^{\frac{U_{\text{eff}}}{T_{\text{eff}}}} = e^{\frac{(1+\tau k)}{D} k \frac{x^2}{2}}, \quad (24)$$

which corresponds to equation (2) where all terms of order τ^2 are zero.

- (b) The **fluctuation path** is obtained using the time reversal symmetry of the problem, namely, by reversing the friction term in the relaxation (noiseless) equations

$$\begin{cases} \dot{x} = p, \\ \dot{p} = +\gamma p - \nabla U_{\text{eff}}, \end{cases}$$

which equivalently gives

$$\ddot{x} - \left(k + \frac{1}{\tau}\right) \dot{x} + \frac{k}{\tau} x = 0. \quad (25)$$

equation (25) together with the boundary conditions $x(0) = x$ and $\dot{x}(0) = 0$ gives

$$x(t) = \frac{x}{1 - k\tau} \left(e^{kt} - k\tau e^{\frac{t}{\tau}} \right). \quad (26)$$

The ‘bulk’ contribution (in the regime $|t| \gg \tau$) and the boundary layer contribution (in the regime $|t| \sim \tau$) are easily identified in equation (26) as the two exponential terms. Figure 1 displays the fluctuation path (26) for decreasing values of τ . It is clearly seen that the path comes closer to the equilibrium fluctuation path of equation $\dot{x} = kx$, when τ decreases toward 0. However, the fluctuation path always has to satisfy the boundary condition $\dot{x}(0) = 0$: this constraint is responsible for the existence of the boundary layer of a size of order τ , and creates a singularity at $t = 0$ in the fluctuation path in the limit $\tau \rightarrow 0$. This surprising phenomenon explains the breakdown of the standard perturbative expansion of the fluctuation path, and creates, as we show, non-trivial local contributions to the quasipotential equation (2) to all orders in τ .




Figure 1. Fluctuation path for the AOUP in a harmonic potential $U(x) = \frac{x^2}{2}$. **Left:** the graph displays the fluctuation paths in the phase space (x, v) , for the three values $\tau = 1.0/0.2/0.05$. The light blue arrows represent the vector field for $\tau = 0.5$. The dark blue line gives the conditions for spatial stationary state $\dot{x} = 0$, and the red line is the equation $v = 2U'(x)$ for the equilibrium fluctuation path (i.e. $\tau = 0$). It can be seen that the convergence to the equilibrium path in the limit $\tau \rightarrow 0$ displays a singularity close to $x(0) = 1.5$. **Right:** the trajectories $\dot{x}(t)$ for the same values of τ as in the left picture. The fluctuation path satisfies $\dot{x}(0) = 0$, whatever the value of τ . The boundary layer of size $\propto \tau$ can clearly be identified close to $t = 0$.

2.2.2. A ratchet. We next illustrate that a consequence of the non-local contribution to the quasipotential (2) is a ratchet effect. We consider the AOUP in the domain $[-1, 1]$ with reflecting boundary conditions (walls) at $x = \pm 1$. We choose the continuous asymmetric potential

$$U(x) = \begin{cases} 1 - 3x^2 - 2x^3 & \text{for } x \in [-1, 0], \\ 1 - 6x^2 + 8x^3 - 3x^4 & \text{for } x \in [0, 1]. \end{cases}$$

The function $U(x)$ is the blue curve displayed in figure 2. For this problem, we compute, using equation (2), the quasipotential Φ_τ defined by the relation

$$D \log \rho(x) = \Phi_\tau(x) + C,$$

where C is some additional constant. Figure 2 displays the function $\Phi_\tau(x)$ truncated respectively to orders τ^0 , τ and τ^2 . It is shown that to order τ^2 , there is an offset $\Delta\Phi$ given by

$$\Delta\Phi = \Phi_\tau(-1) - \Phi_\tau(1) = -\tau^2 \int_{-1}^1 |U'(y)|^2 U'''(y) dy.$$

The offset is responsible for the accumulation of active particles on the right side of the ratchet potential, according to the relation $\frac{\rho(1)}{\rho(-1)} \underset{D \rightarrow 0}{\asymp} e^{\frac{\Delta\Phi}{D}}$. This effect is absent if one considers the expansion of Φ_τ up to order τ only.




Figure 2. The ratchet effect for AOUP. The blue curve is the ratchet potential $U(x)$, which also corresponds to the quasipotential at equilibrium $\Phi_{\tau=0}$. The figure displays the quasipotential Φ_τ truncated to order τ (yellow curve), and τ^2 (red curve), for $\tau = 0.1$. It can be seen that an offset $\Delta\Phi$ appears between the right and the left of the ratchet to order τ^2 , creating an accumulation of active particles on the right of the ratchet.

2.3. The large τ limit

We now turn to discuss the large τ limit. In particular, we illustrate how results can be obtained using heuristic arguments. The full details of the calculation are presented in section 4. The limit is best understood by rescaling time according to $t \leftarrow t/\tau$ and $D \leftarrow D/\tau$ in equation (1). This yields

$$\begin{cases} \dot{x} = \tau(v - U'(x)), \\ \dot{v} = -v + \sqrt{2D}\xi(t). \end{cases} \quad (27)$$

From this it is easy to see that in the large τ limit the dynamical relaxation time in the trap is much smaller than the time scale associated with the variation of the active force. In particular, in the strict limit $\tau \rightarrow \infty$ the particle is in a quasistationary state given by the equation

$$U'(x(t)) = v(t). \quad (28)$$

Since the dynamics of v does not depend on x , the location of the particle can be determined from equation (28) by inverting the function U' . Figure 3 displays the dynamics in phase space (x, v) for $\tau = 30$. This implies that, for a typical barrier, the curve $v = U'(x)$ has a stable manifold in an interval $[x_0, x_{\text{cr}}]$ and an unstable manifold in $[x_{\text{cr}}, x_s]$. Here x_{cr} is the (unique) inflection point at which $U''(x_{\text{cr}}) = 0$.

This implies that in order to cross the barrier, the particle has to reach x_{cr} by moving along the stable manifold, and then, once x_{cr} is reached, it can ‘fly over’ the barrier using the deterministic dynamics. Namely, any position $x > x_{\text{cr}}$ can be reached without any additional action cost, once the inflection point x_{cr} has been crossed. We note that this phenomenology seems to be generic for fluctuation paths in the presence of a slow manifold as shown in [26].

The barrier displayed in figure 3 is thus a **force barrier** and not an energy barrier as found in equilibrium problems. Then the mean escape time in the large deviation

Nonlocal stationary probability distributions and escape rates for an active Ornstein–Uhlenbeck particle




Figure 3. (a) Phase space structure of AOUP in the large τ limit. The dynamics along x is very fast compared to the v direction. The blue arrows represent the deterministic dynamics for $\tau = 30$, and $U(x) = \frac{x^2}{2} - \frac{x^3}{3}$. To escape from the metastable state, the process first follows the stable manifold up to x_{cr} and then can fly over the barrier using the deterministic field. (b) Graph of the potential barrier $U(x)$. The saddle point is $x_s = 1$ and the inflection point is $x_{\text{cr}} = 0.5$.

regime $D \rightarrow 0$ is given by the mean time required for the Ornstein–Uhlenbeck process $v(t)$ to reach $F_{\text{max}} = -U'(x_{\text{cr}})$. The result is

$$\langle T_{\text{esc}} \rangle \underset{D \rightarrow 0}{\asymp} e^{\frac{|F_{\text{max}}|^2}{2D}}. \quad (29)$$

Following the same line of arguments, one can obtain the quasipotential in the large τ limit

$$\Phi_\tau(x) \xrightarrow{\tau \rightarrow +\infty} \begin{cases} \frac{1}{2} |U'(x)|^2 & \text{for } x_0 < x < x_{\text{cr}}, \\ \frac{1}{2} |U'(x_{\text{cri}})|^2 & \text{for } x_{\text{cr}} < x < x_s. \end{cases} \quad (30)$$

Whereas the result of equation (30) is relatively easy to obtain, the first order correction is not. One has to deal with two different types of boundary layers in the fluctuation path, depending on the value of x .

- For $x < x_{\text{cr}}$, the fluctuation path has a boundary layer of size $\frac{1}{\tau}$ close to $t = 0$. The expansion of Φ_τ can be done in powers of $\frac{1}{\tau}$, and one obtains to leading order

$$\Phi_\tau(x) \underset{\tau \rightarrow +\infty}{\sim} \frac{|U'(x)|^2}{2} + \frac{|U'(x)|^2}{2\tau U''(x)} + O(1/\tau^2). \quad (31)$$

Note that as required this expression agrees with the exact expression (24) obtained for a harmonic potential.

$x > x_2$ is in fact the same as to reach $x > x_1$. The fraction of time spent in the second well $x_1 < x < x_2$ vanishes in the limit $\tau \rightarrow +\infty$. To show this latter point, we implemented a numerical simulation where we record the total times T_1 and T_2 spent by each particle in the first and second well respectively, and we numerically compute the ratio

$$\langle r \rangle = \left\langle \frac{T_2}{T_1 + T_2} \right\rangle, \quad (33)$$

where the average is done over escape events, for each value of τ . Simple scaling arguments show that the large τ behavior of $\langle r \rangle$ is

$$\langle r(\tau) \rangle \underset{\tau \rightarrow +\infty}{\propto} \frac{\log(\tau)}{\tau}.$$

We show in figure 4 that the function $\langle r \rangle$ indeed decreases slowly with τ . In addition, we illustrate the result with the two supplementary movies.

3. AOUP in the small τ limit

We now detail the full calculation of the quasipotential in the small τ limit. Here we use a Lagrangian approach described in section 2.1, that allows for a generic organization of the perturbation expansion. The Lagrangian is given in equation (8). Recall that the quasipotential at a point $x \in \mathcal{B}_0$ is obtained by evaluating the action on the ‘instanton trajectory’ whose solution is given through the saddle-point equation

$$\ddot{x} - U'(x)U''(x) - \tau^2 (\ddot{\ddot{x}} + \dot{x}^3 U^{(4)}(x) - \ddot{x} (U''(x)^2 - 3\dot{x}U'''(x)) - \dot{x}^2 U'''(x)U''(x)) = 0, \quad (34)$$

Assuming that x_0 is the location of the optimal minimum (the generalization being straightforward using standard methods, see for example [34]) the optimal path $x(t)$ starts from the local minimum x_0 of U in \mathcal{B}_0 at time $t = -\infty$ and arrives in x at time $t = 0$. The Hamiltonian boundary condition (14) on the momentum P_v vanishing at $t = 0$ implies that $\dot{x}(0) = 0$. The boundary conditions on equation (34) are given in equation (9). Finally, note that conservation of ‘energy’ [see equations (12) and (16)] enforces the optimal path to satisfy

$$H(x, \dot{x}, \ddot{x}, \ddot{\ddot{x}}) \equiv -U'(x)^2 + \tau^2 (-2\dot{x}^3 U'''(x) + \dot{x}^2 U''(x)^2 + \ddot{x}^2 - 2\dot{x}\ddot{x}) + \dot{x}^2 = 0. \quad (35)$$

3.1. Expansion order by order

As shown in section 2.2.1 (see figure 1) for the harmonic case, the instanton solution $x(t)$ presents two distinct regimes of time in the small τ limit: (i) a ‘bulk’ regime $t \ll -\tau$ where the solution is uniformly close to the $\tau = 0$ instanton, and where an expansion in powers of τ is well defined; and (ii) a ‘boundary-layer’ regime $t \sim -\tau$ where the instanton cannot be obtained as a small perturbation around the $\tau = 0$ instanton solution. One

therefore has to treat the boundary layer using a singular perturbation theory. To this end, in what follows we determine order by order in powers of τ the form of the instanton in the bulk and in the boundary layer; these forms are determined up to constants that we obtain using standard asymptotic matching techniques.

Namely, we write the instanton solution in the bulk regime as

$$x(t) = x_B(t) \equiv x^{(0)}(t) + \tau x^{(1)}(t) + \tau^2 x^{(2)}(t) + \tau^3 x^{(3)}(t) + \dots, \quad (36)$$

with the boundary condition $x_B(-\infty) = x_0$. In the boundary layer, following standard approaches, we introduce $x(t) = x_{bl}(t/\tau)$ and defines $t' = t/\tau$. The solution in this regime is expanded as

$$x_{bl}(t') = x_{bl}^{(0)}(t') + \tau x_{bl}^{(1)}(t') + \tau^2 x_{bl}^{(2)}(t') + \tau^3 x_{bl}^{(3)}(t') + \dots, \quad (37)$$

and equation (9) implies that $x_{bl}(0) = 0$ and $\dot{x}_{bl}(0) = 0$. The bulk and boundary solution can be asymptotically matched by demanding that $x_B(t)|_{t \rightarrow 0}$ and $x_{bl}(t/\tau)|_{t \rightarrow -\infty}$ have the same functional form. The overall structure of the perturbation theory is implemented by enforcing the zero-energy condition (35) order by order in τ for both expansions for the bulk (36) and the boundary layer (37).

In what follows we specify the solution order by order.

3.1.1. Zeroth order. In the bulk regime, using (36) with (35) one finds at *zeroth* order in τ that

$$(U'(x^{(0)}))^2 - (\dot{x}^{(0)})^2 = 0,$$

One thus sees that $x^{(0)}(t)$ is equal to the equilibrium instanton $\tilde{x}(t)$, which is given by the solution of

$$\dot{\tilde{x}}(t) = U'(\tilde{x}(t)) \quad \text{with } \tilde{x}(-\infty) = x_0 \quad \text{and} \quad \tilde{x}(0) = x. \quad (38)$$

Next, we express higher orders of the bulk expansion (36) as a function of $\tilde{x}(t)$.

In the boundary-layer regime, using equations (35) and (37) one finds to *zeroth* order in τ that

$$2\dot{x}_{bl}^{(0)}\ddot{x}_{bl}^{(0)} - \left(\ddot{x}_{bl}^{(0)}\right)^2 - \left(\dot{x}_{bl}^{(0)}\right)^2 = 0,$$

whose solution is

$$x_{bl}^{(0)}(t') = \left(2C_1 t' - C_1^2 e^{-t'} + e^{t'}\right) C_2 + C_3,$$

with C_k 's constants. The asymptotic matching condition implies that $x_{bl}^{(0)}(t')$ has to remain constant as $t' \rightarrow -\infty$. This imposes $C_2 = 0$, which with the boundary condition at $t' = 0$ implies $C_3 = x$. Therefore, $x_{bl}^{(0)}(t') = x$.

The quasipotential is obtained by decomposing the time interval $]-\infty, 0]$ into $]-\infty, -A\tau] \cup]-A\tau, 0]$ and using the bulk and boundary-layer expressions of the instanton trajectory:

$$\Phi(x) = \int_{-\infty}^{-A\tau} \mathcal{L}(x_B, \dot{x}_B) dt + \int_{-A}^0 \tau \mathcal{L}(x_{bl}, \tau \dot{x}_{bl}) dt', \quad (39)$$

Clearly, for this to be consistent the result should be independent of A in the large A limit. Using the solution we have obtained one finds, as expected, that the boundary layer does not contribute at order τ^0 and one recovers the equilibrium result

$$\Phi(x) = U(x) - U(x_0) + O(\tau).$$

3.1.2. First order. In the bulk regime, using (35) and (36) and taking into account zeroth order results, one finds that

$$\dot{\tilde{x}}\dot{x}^{(1)} - \ddot{\tilde{x}}x^{(1)} = 0,$$

where we also used equation (38) for the equilibrium instanton $\tilde{x}(t)$, to eliminate derivatives of U . The solution is then

$$x^{(1)}(t) = c_1 \dot{\tilde{x}}(t). \quad (40)$$

In the boundary-layer regime, equations (34) and (37) give

$$\ddot{x}_{\text{bl}}^{(1)} - \ddot{\tilde{x}}_{\text{bl}}^{(1)} = 0,$$

whose solution, satisfying the boundary conditions⁴ at $t' = 0$ is

$$x_{\text{bl}}^{(1)}(t') = D_1 (e^{t'} - 1 - t') + D_2 (e^{-t'} - 1 + t')$$

and the zero-energy condition, together with the solution at zeroth order, imposes $(D_1 + D_2)^2 - U'(x)^2 = 0$.

The asymptotic matching then imposes $D_2 = 0$ so that $D_1 = \pm U'(x)$. To determine the sign of D_1 and the value of c_1 in (40) one compares the asymptotic behaviors of the bulk and the boundary-layer:

$$\begin{aligned} x_{\text{B}}(t)|_{t=\tau t'} &\underset{t \rightarrow 0}{\approx} x + [c_1 U'(x) + t' U'(x)] \tau + \dots \\ x_{\text{bl}}(t') &\underset{t' \rightarrow -\infty}{\approx} x - [D_1(1 + t')] \tau + \dots \end{aligned}$$

(Note that for $x_{\text{B}}(t)$ both $x^{(0)}(t)$ and $x^{(1)}(t)$ contribute, and we used the equation of motion $\dot{\tilde{x}}(t) = U'(\tilde{x}(t))$ to evaluate $\dot{\tilde{x}}(0) = U'(x)$.) The asymptotic matching therefore imposes $D_1 = -U'(x)$ and $c_1 = 1$. To summarize, the first-order perturbations in the bulk and boundary layer are:

$$x^{(1)}(t) = \dot{\tilde{x}}(t); \quad x_{\text{bl}}^{(1)}(t') = (e^{t'} - 1 - t') U'(x). \quad (41)$$

We remark that at this first order, the boundary layer $x_{\text{bl}}(t/\tau)$ contains terms proportional to $\tau e^{t/\tau}$ which are non-perturbative in τ as $\tau \rightarrow 0$ at fixed t . This fact explains how the boundary layer contributes to the expression of the quasipotential in $\Phi(x) = \int_{-\infty}^0 \mathcal{L}(x(t), \dot{x}(t)) dt$ even though it only affects this time integral over a duration τ .

⁴ Note that since $x_{\text{bl}}(0) = x$ the boundary condition on the higher order terms in $t' = 0$ are $x_{\text{bl}}^{(k)}(0) = 0$ and $\dot{x}_{\text{bl}}^{(k)}(0) = 0$ ($k \geq 1$).

As we discussed before equation (39) the quasipotential is given by

$$\Phi(x) = \lim_{A \rightarrow \infty} [\Phi_B^A(x) + \Phi_{bl}^A(x)] \quad (42)$$

with

$$\Phi_B^A(x) = \int_{-\infty}^{-A\tau} \mathcal{L}(x_B, \dot{x}_B) dt \quad \text{and} \quad \Phi_{bl}^A(x) = \int_{-A}^0 \tau \mathcal{L}(x_{bl}, \tau \dot{x}_{bl}) dt'. \quad (43)$$

Using the solution for $x_B(t)$ one finds to order τ

$$\mathcal{L}(x_B, \dot{x}_B) = \dot{x}U'(\tilde{x}) + 4\tau\dot{x}\ddot{x} = \partial_t [U(\tilde{x}) + 2\tau\dot{x}^2],$$

so that

$$\Phi_B^A(x) = [U(\tilde{x}) + 2\tau\dot{x}^2]_{-\infty}^{-A\tau} = [U(\tilde{x}) + 2\tau U'(\tilde{x})^2]_{-\infty}^{-A\tau}.$$

Using then $U'(\tilde{x}(-\infty)) = U'(x_0) = 0$ and $\tilde{x}(-A\tau) = \tilde{x}(0) - A\tau\dot{x}(0) = \tilde{x}(0) - A\tau U'(x)$ one finally has

$$\Phi_B^A(x) = U(x) - U(x_0) + 2\tau(-A + 1)U'(x)^2 + O(\tau^2). \quad (44)$$

In the boundary layer one finds, up to order τ

$$\tau \mathcal{L}(x_{bl}, \tau \dot{x}_{bl}) = \tau (e^{t'} - 1)^2 U'(x)^2 + O(\tau^2)$$

so that equation (43) gives

$$\Phi_{bl}^A(x) = \frac{1}{2}\tau [2A - 3 - e^{-2A} + 4e^{-A}] U'(x)^2 + O(\tau^2). \quad (45)$$

Adding (44) and (45) one sees that the terms linear in A compensate as we take $A \rightarrow \infty$. We finally obtain from (42) the quasipotential to order τ

$$\Phi(x) = U(x) - U(x_0) + \frac{1}{2}\tau U'(x)^2 + O(\tau^2). \quad (46)$$

Remark. If we had ignored the boundary layer, we would obtain that $c_1 = 0$ in (40) so as to satisfy the boundary condition (9) in $t = 0$ [because $\dot{x}(0) = U'(x) \neq 0$ in general], meaning that the instanton $x(t)$ would present no correction of order τ . This would imply that the only contribution of order τ to the quasipotential is given by the double-product in the Lagrangian (8), that is:

$$\frac{1}{4}\tau \int \partial_t [(\dot{x} + U'(\tilde{x}))^2] dt = \frac{1}{4}\tau [(\dot{x} + U'(\tilde{x}))^2]_{-\infty}^0 = \tau U'(x)^2$$

which is wrong by a factor $\frac{1}{2}$ compared to the actual result (46). This illustrates how the presence of the boundary layer affects the determination of the quasipotential: it modifies the boundary condition in $t \rightarrow 0$ seen by the instanton, and it contributes at order τ (and higher orders) to the integral of the Lagrangian in the action.

3.1.3. Second order. In the bulk regime, inserting equation (36) in the zero-energy condition (35) and taking into account the result obtained at *zeroth* and first order, one finds that the second-order correction to the bulk instanton is given by the solution of

$$\dot{\tilde{x}}\dot{\tilde{x}}^{(2)} - \ddot{\tilde{x}}\tilde{x}^{(2)} + 2(\ddot{\tilde{x}}^2 - \dot{\tilde{x}}\ddot{\tilde{x}}) = 0.$$

Here we used equation (38) to eliminate occurrences of $U(\tilde{x})$ and its derivatives. The solution to this equation is given by

$$x^{(2)}(t) = d_1 \dot{\tilde{x}}(t) + \frac{5}{2} \ddot{\tilde{x}}(t),$$

where d_1 is a constant. In the boundary-layer regime, inserting equation (37) in (34) and using the results obtained at *zeroth* and first order one finds in a similar manner

$$\ddot{\tilde{x}}_{\text{bl}}^{(2)} - \ddot{\tilde{x}}_{\text{bl}}^{(2)} + U'(x)U''(x) = 0.$$

Using the boundary conditions $x_{\text{bl}}^{(2)}(0) = 0$ and $\dot{x}_{\text{bl}}^{(2)}(0) = 0$ with the zero-energy constraint and demanding that the solution does not diverge exponentially as $t \rightarrow -\infty$, so that the bulk solution can be matched, gives

$$x_{\text{bl}}^{(2)}(t') = \left(1 + t' + \frac{t'^2}{2} - e^{t'}\right) U'(x)U''(x). \quad (47)$$

Then using asymptotic matching one finds $d_1 = -\frac{3}{2}U''(x)$ so that finally

$$x_{\text{B}}^{(2)}(t) = -\frac{3}{2}U''(x)\dot{\tilde{x}}(t) + \frac{5}{2}\ddot{\tilde{x}}(t). \quad (48)$$

We now proceed to evaluate the quasipotential. The order τ calculation of the quasipotential gave a local function of the potential, see equation (46). We now show that at order τ^2 the quasipotential, as stressed in the introduction, becomes non-local.

We saw that the solution to order τ for the functions $\Phi_{\text{B}}^A(x)$ and $\Phi_{\text{bl}}^A(x)$ had divergent contributions at order τ when taking the $A \rightarrow \infty$ limit which compensated each other when $\Phi_{\text{B}}^A(x)$ and $\Phi_{\text{bl}}^A(x)$ were summed. Performing a similar analysis at order τ^2 (and higher) is more complex. Here we avoid this procedure. To do so we replace $\mathcal{L}(x, \dot{x})$ in the integrals of equation (43) defining $\Phi_{\text{B}}^A(x)$ and $\Phi_{\text{bl}}^A(x)$ by

$$\mathcal{L}_F(x, \dot{x}) = \mathcal{L}(x, \dot{x}) - \partial_t \underbrace{[F_0(x) + \tau F_1(x) + \dots]}_{\equiv F(x)}. \quad (49)$$

Namely, we remove a counter-term that is a total derivative. In the action, the counter-term becomes a boundary term that can be evaluated separately, before any expansion (and before splitting $]-\infty, 0]$ into a bulk and boundary-layer regime). In sum, we now write

$$\Phi(x) = [F(x(t))]_{x_0}^x + \lim_{A \rightarrow \infty} [\Phi_{\text{B}}^A(x) + \Phi_{\text{bl}}^A(x)], \quad (50)$$

with

$$\Phi_B^A(x) = \int_{-\infty}^{-A\tau} \mathcal{L}_F(x_B, \dot{x}_B) dt \quad \text{and} \quad \Phi_{bl}^A(x) = \int_{-A}^0 \tau \mathcal{L}_F(x_{bl}, \tau \dot{x}_{bl}) dt'. \quad (51)$$

The functions $F_0(x)$ and $F_1(x)$ in equation (49) are taken so that asymptotic matching does not give any divergence as $A \rightarrow \infty$. Taking

$$F_0(x) = U(x) \quad \text{and} \quad F_1(x) = U'(x)^2, \quad (52)$$

one finds after a straightforward expansion of the integrals in (51), separately, and using (50) that, as announced previously

$$\Phi(x) = U(x) - U(x_0) + \frac{1}{2} \tau U'(x)^2 - \frac{1}{2} \tau^2 \int_{x_0}^x [U'(y)]^2 U'''(y) dy + O(\tau^3). \quad (53)$$

For instance, the choice of function for $F_0(x)$ avoids the appearance of terms linear in A in the bulk and boundary-layer expressions (44) and (45) to order τ . At order τ^2 one finds by inserting the series (37) into (51) that the contribution of order τ^2 to the integrand of (51) is

$$\begin{aligned} \tau \mathcal{L}_F(x_{bl}, \tau \dot{x}_{bl})|_{\tau^2} = & \left\{ 3U'(x)^2 U''(x) e^{2t'} + U'(x) [F_1'(x) - (t' + 5) U'(x) U''(x)] e^{t'} \right. \\ & \left. + U'(x) [2U'(x) U''(x) - F_1'(x)] \right\} \tau^2. \end{aligned} \quad (54)$$

When performing the integral over t' in (51), the second line in (54) yields terms which diverge with $A \rightarrow \infty$. The choice of function $F_1(x)$ in equation (52) then avoids this divergence.

We note that the result (53), derived here through a singular perturbation theory for the instanton, can also be obtained through a different approach. In an early work, Kłosek-Dygas *et al* [3] studied the escape problem using a Fokker–Planck approach (and in the small τ expansion). The approach yields the steady-state solution $P_{st}(x, \dot{x})$ of the Fokker–Planck equation on the joint probability density $P(x, \dot{x}, t)$, within a large-deviation approach at small D . The authors obtain an expansion of $\lim_{D \rightarrow 0} D \log P_{st}(x, \dot{x})$ in powers of τ (their equation (21)). If one takes the marginal over the variable \dot{x} in their results it is easy to check that the result is again (53). This alternative derivation however does not unveil the singular nature of the optimal trajectory.

4. AOUP in the large τ limit

In this part it is more convenient to use the Hamiltonian formalism. The large τ limit is best understood by rescaling time according to $t \leftarrow t/\tau$. After the time rescaling, the Hamiltonian (12) takes the form

$$H(x, v, P_x, P_v) = \tau (v - U'(x)) P_x - v P_v + P_v^2, \quad (55)$$

so that the instanton equations are

$$\begin{cases} \dot{x} = \tau(v - U'(x)), \\ \dot{v} = -v + 2P_v, \end{cases} \quad (56)$$

and

$$\begin{cases} \dot{P}_x = \tau U''(x)P_x, \\ \dot{P}_v = P_v - \tau P_x. \end{cases} \quad (57)$$

For ease of presentation, we now split the discussion in the escape problem and the calculation of the general quasipotential. In both cases it will be useful to keep in mind a general potential of the form presented in figure 3.

4.1. The quasipotential for $x > x_{\text{cr}}$

As before we denote the position of the highest point of the potential by x_s and the position of the (unique) inflection point where $U''(x_{\text{cr}}) = 0$ by x_{cr} . To evaluate the quasipotential we look for an *instanton path* by solving equations (56) and (57) with the boundary conditions

$$x(t), v(t) \xrightarrow{t \rightarrow -\infty} 0; \quad P_x(t), P_v(t) \xrightarrow{t \rightarrow -\infty} 0. \quad (58)$$

The boundary conditions at later times are more subtle. Initially using the time translation invariance of the instanton path we choose

$$x(0) = x_{\text{cr}}, \quad (59)$$

but, as will become clear, we also have to specify boundary conditions at positive times. This will be carried out below.

4.1.1. Bulk solution. We first show that $P_x = 0$ to any order for t not close to $t = 0$, where as we will show, a boundary layer appears. Consider $\delta > 0$, and the instanton path $x(t)$ on the interval $]-\infty, -\delta]$: as $x(t)$ is a monotonic function of t , it is clear that $x(t) < x(-\delta) < 0$ on this part of the path. Moreover, since we assume the existence of a single inflection point, U'' is strictly positive in $[x_0, x(-\delta)]$. Therefore, there exists some constant $c_\delta > 0$ such that

$$\forall t < -\delta, \quad U''(x(t)) > c_\delta. \quad (60)$$

Then the first equation of (57) shows that $\dot{P}_x > c_\delta \tau P_x$ over the time interval $]-\infty, -\delta]$. Thus, we deduce the bound

$$\forall t < -\delta, \quad |P_x(t)| < |P_x(-\delta)| e^{c_\delta \tau t}.$$

The momentum P_x relaxes faster than $e^{c_\delta \tau t}$ when $t \rightarrow -\infty$. A similar argument shows that P_x relaxes exponentially faster than $e^{c'_\delta \tau t}$, with $c'_\delta < 0$ when $t \rightarrow +\infty$. Therefore, since we are interested in the limit $\tau \rightarrow \infty$, these bounds prove that an expansion of P_x in powers of τ vanishes to any order in τ , except in the immediate vicinity of $x_{\text{cr}} = 0$.

This simple observation indicates the existence of a boundary layer in the vicinity of $t = 0$ that connects the part $]-\infty, 0[$ to the part $]0, +\infty[$ of the instanton path. The scaling with τ of the boundary layer is nontrivial, and will be derived in the following.

In this section, we solve equations (56) and (57) in the bulk regimes $t < 0$ and $t > 0$. The boundary layer (t close to zero) is solved in section 4.1.2. With $P_x = 0$, the instanton equations in the bulk are

$$\begin{cases} \dot{x} = \tau (v - U'(x)), \\ \dot{v} = -v + 2P_v, \\ \dot{P}_v = P_v. \end{cases} \quad (61)$$

To solve the system (61), we expand the solution as

$$P_v(t) = P_v^{(0)}(t) + \frac{1}{\tau^n} P_v^{(1)}(t) + \frac{1}{\tau^{2n}} P_v^{(2)}(t) + \dots, \quad (62)$$

and similarly for $x(t)$ and $v(t)$. The exponent n is related to the size of the boundary layer close to $t = 0$, and has to be determined self-consistently. However, before considering the corrections we first solve the *zeroth* order of the problem.

Zeroth order: to this order we have, using equation (61)

$$\begin{cases} U'(x^{(0)}) = v^{(0)}, \\ \dot{v}^{(0)} = -v^{(0)} + 2P_v^{(0)}, \\ \dot{P}_v^{(0)} = P_v^{(0)}. \end{cases} \quad (63)$$

The solution for $P_v^{(0)}$ and $v^{(0)}$ is straightforward. Taking into account the boundary conditions (58) in $t = -\infty$, we get

$$t < 0: \begin{cases} v^{(0)}(t) = A_0 e^t, \\ P_v^{(0)}(t) = A_0 e^t \end{cases} \quad \text{and} \quad t > 0: \begin{cases} v^{(0)}(t) = B_0 e^{-t}, \\ P_v^{(0)}(t) = 0. \end{cases} \quad (64)$$

where $A_0 > 0$. Since the third equation in (63) implies that P_v is either zero identically or diverges to infinity for $t > 0$ one has to take $P_v = 0$ as expected. Beyond x_{cr} the motion of the instanton follows a noiseless relaxation path. Therefore its contribution to the action is zero. Then the solution for $v^{(0)}$ follows with $B_0 > 0$.

The solution for $x^{(0)}(t)$ has to be computed from the first equation of (63), inverting the function U' . In the vicinity of $x_{\text{cr}} = 0$, we can expand U' as

$$U'(x) = \beta - \gamma x^2 + O(x^3).$$

Then using $U'(x^{(0)}) = v^{(0)}$, and the solution (64) we get

$$x^{(0)}(t) \underset{t \rightarrow 0^-}{\sim} \sqrt{\frac{1}{\gamma} (\beta - A_0 e^t)}.$$

Note that since the instanton path for $t > 0$ does not contribute to the action we do not have to consider the solution in that regime. The constraint $x(0) = 0$ imposes $A_0 = \beta$ and thus

$$x^{(0)}(t) \underset{t \rightarrow 0^-}{\sim} -\sqrt{\frac{\beta}{\gamma}}|t| \quad (65)$$

equation (65) gives the asymptotic behavior that the boundary layer solution has to satisfy. As we show just below, it also constrains the size of the boundary layer.

Boundary layer scaling: in the vicinity of $x_{\text{cr}} = 0$, the first equation of (57) gives

$$\dot{P}_x = -2\tau\gamma x P_x. \quad (66)$$

Plugging the solution (65) into (66) gives

$$\dot{P}_x = -2\tau\sqrt{\gamma\beta}|t|P_x,$$

which can be integrated to give

$$P_x(t) \propto e^{-2\sqrt{\gamma\beta}\frac{2\tau}{3}|t|^{3/2}}.$$

This proves that P_x relaxes over a typical time scaling $\propto \frac{1}{\tau^{2/3}}$. The boundary layer scaling variable is also $\tau^{2/3}t$, as we show below. In what follows it will become clear that this implies that for the expansion (62) to be consistent one has to take $n = \frac{1}{3}$.

Higher orders: using the linearity of the bulk equations for v and P_v , we can express the solution to any order k as

$$t < 0: \begin{cases} v^{(k)}(t) = A_k e^t, \\ P_v^{(k)}(t) = A_k e^t \end{cases} \quad \text{and} \quad t > 0: \begin{cases} v^{(k)}(t) = B_k e^{-t}, \\ P_v^{(k)}(t) = 0. \end{cases} \quad (67)$$

One can then find the solution $x^{(k)}(t)$ using the expansion of $\dot{x} = \tau(v - U'(x))$ to k th order in $\frac{1}{\tau^{1/3}}$. The constants A_k and B_k have to be determined by matching the bulk solution to the boundary layer as we do below to up to order two. We now turn to consider the solution in the boundary layer in the vicinity of x_{cr} .

4.1.2. Boundary layer solution. To find the boundary layer equations, we take equations (56) and (57) and again use the expansion of $U'(x)$ to second order near $x_{\text{cr}} = 0$

$$U'(x) = \beta - \gamma x^2 + O(x^3).$$

This gives

$$\begin{cases} \dot{x} = \tau(v - \beta + \gamma x^2), \\ \dot{v} = -v + 2P_v, \end{cases} \quad (68)$$

and

$$\begin{cases} \dot{P}_x = -2\tau\gamma x P_x, \\ \dot{P}_v = P_v - \tau P_x. \end{cases} \quad (69)$$

Using the two parameters β and γ , we can rescale all fields in order to arrive at non-dimensional equations. We set $\tau' = \sqrt{\gamma\beta}\tau$, $v' = v/\beta$, $P_v' = P_v/\beta$, $x' = x\sqrt{\frac{\gamma}{\beta}}$ and $P_x' = \tau P_x/\beta$. We get

$$\begin{cases} \dot{x}' = \tau' (v' - 1 + x'^2), \\ \dot{v}' = -v' + 2P_v', \end{cases} \quad (70)$$

and

$$\begin{cases} \dot{P}_x' = -2\tau' x' P_x', \\ \dot{P}_v' = P_v' - P_x'. \end{cases} \quad (71)$$

Finally, we use the scaling of the boundary layer and we rescale time according to $t' = \tau'^{2/3}t$ and for self-consistency define the variable $x'' = x'\tau'^{1/3}$ and $P_x'' = P_x'/\tau'^{2/3}$. Omitting the primes, the boundary layer equations are

$$\begin{cases} \dot{x} = \tau^{2/3} (v - 1) + x^2, \\ \dot{v} = -\frac{1}{\tau^{2/3}}v + \frac{2}{\tau^{2/3}}P_v, \end{cases} \quad (72)$$

and

$$\begin{cases} \dot{P}_x = -2xP_x, \\ \dot{P}_v = \frac{1}{\tau^{2/3}}P_v - P_x. \end{cases} \quad (73)$$

We then expand the boundary layer solution as

$$x_{\text{bl}}(t) = x_{\text{bl}}^{(0)}(t) + \frac{1}{\tau^{1/3}}x_{\text{bl}}^{(1)}(t) + \frac{1}{\tau^{2/3}}x_{\text{bl}}^{(2)}(t) + \dots$$

and similar expansions for $P_{x,\text{bl}}$, $P_{v,\text{bl}}$, $v_{\text{bl}}(t)$. We now turn to solve these equation order by order.

Leading order: in this case the equations are dominated by terms of order $\tau^{2/3}$. This gives

$$v_{\text{bl}}^{(0)} = 1,$$

which is consistent with the bulk *zeroth* order solution found in (64) with $A_0 = \beta$ (see the discussion before equation (65)). This also implies $B_0 = \beta$. It is then easy to see that the next term (of order $\tau^{1/3}$) vanishes:

$$v_{\text{bl}}^{(1)} = 0,$$

which implies $A_1 = 0$ and $B_1 = 0$ in the bulk, and that all terms of order $\frac{1}{\tau^{1/3}}$ vanish in the bulk solution (67).

Zeroth order: to next order the equation are

$$\begin{cases} \dot{x}_{\text{bl}}^{(0)} = v_{\text{bl}}^{(2)} + \left(x_{\text{bl}}^{(0)}\right)^2, \\ \dot{v}_{\text{bl}}^{(2)} = -1 + 2P_{v,\text{bl}}^{(0)}, \end{cases} \quad (74)$$

and

$$\begin{cases} \dot{P}_{x,\text{bl}}^{(0)} = -2x_{\text{bl}}^{(0)} P_{x,\text{bl}}^{(0)}, \\ \dot{P}_{v,\text{bl}}^{(0)} = -P_{x,\text{bl}}^{(0)}. \end{cases} \quad (75)$$

The boundary conditions for the set of equations (74) and (75) can be found by matching them with the bulk solution. The fact that $P_x = 0$ to any order in the bulk gives

$$P_{x,\text{bl}}^{(0)} \xrightarrow{t \rightarrow \pm\infty} 0. \quad (76)$$

The matching with the bulk solution (64) with $A_0 = B_0 = \beta$ gives

$$\begin{cases} P_{v,\text{bl}}^{(0)} \xrightarrow{t \rightarrow -\infty} 1, \\ P_{v,\text{bl}}^{(0)} \xrightarrow{t \rightarrow +\infty} 0. \end{cases} \quad (77)$$

The bulk estimate (65) close to $t = 0$ gives

$$x_{\text{bl}}^{(0)}(t) \underset{t \rightarrow -\infty}{\sim} -\sqrt{|t|}, \quad (78)$$

and finally, the instanton path has to cross $x_{\text{cr}} = 0$ at $t = 0$, and then leaves the vicinity of x_{cr} with

$$\begin{cases} x_{\text{bl}}^{(0)}(0) = 0, \\ x_{\text{bl}}^{(0)}(t) \xrightarrow{t \rightarrow +\infty} +\infty. \end{cases} \quad (79)$$

To this order we find a solution which obeys the symmetry

$$\begin{cases} x_{\text{bl}}^{(0)}(t) \leftrightarrow -x_{\text{bl}}^{(0)}(-t), \\ v_{\text{bl}}^{(2)}(t) \leftrightarrow v_{\text{bl}}^{(2)}(-t), \\ P_{v,\text{bl}}^{(0)}(t) \leftrightarrow 1 - P_{v,\text{bl}}^{(0)}(-t), \\ P_{x,\text{bl}}^{(0)}(t) \leftrightarrow P_{x,\text{bl}}^{(0)}(-t). \end{cases} \quad (80)$$

The solution can be computed numerically using an iterative procedure. Figure 5 displays the resulting solutions.

The last step is to match the boundary layer solution at $t \rightarrow -\infty$ to the bulk solution in the limit $t \rightarrow 0$, order by order in τ . In the bulk we have

$$\begin{aligned} v\left(t = \frac{t'}{\tau^{2/3}}\right) &= \left(A_0 + \frac{1}{\tau^{1/3}}A_1 + \frac{1}{\tau^{2/3}}A_2\right) e^{t'/\tau^{2/3}} + O(1/\tau) \\ &\underset{\tau \rightarrow +\infty}{\sim} \left(A_0 + \frac{A_1}{\tau^{1/3}} + \frac{A_2}{\tau^{2/3}}\right) \left(1 + \frac{t'}{\tau^{2/3}}\right) + O(1/\tau), \end{aligned}$$




Figure 5. Numerical resolution of the undimensioned boundary layer equations equations (72) and (73).

and the boundary layer solution $v_{\text{bl}}(t')$ up to second order gives

$$v_{\text{bl}}(t') = \beta v_{\text{bl}}^{(0)}(t) + \frac{1}{\tau^{1/3}} \beta v_{\text{bl}}^{(1)}(t') + \frac{1}{\tau^{2/3}} \beta v_{\text{bl}}^{(2)}(t') + O(1/\tau')$$

$$\underset{t' \rightarrow -\infty}{\sim} \beta + \frac{1}{\tau^{2/3}} \beta (C + t') + O(1/\tau'),$$

where $C > 0$ is some positive constant (see figure 5). Self consistency of the boundary layer expansion and the bulk solution requires $A_0 = \beta$, $A_1 = 0$, and

$$A_2 = \frac{\beta C}{(\sqrt{\gamma\beta})^{2/3}}. \quad (81)$$

C can be expressed in term of the numerical solution of equations (74) and (75). Using the equation for $v_{\text{bl}}^{(2)}(t)$ we have

$$\dot{v}_{\text{bl}}^{(2)} - 1 = 2 \left(P_{v,\text{bl}}^{(0)} - 1 \right).$$

Both sides of the latter equation can be integrated, yielding

$$\int_{-\infty}^0 \frac{d}{dt} \left(v_{\text{bl}}^{(2)} - t \right) dt = 2 \int_{-\infty}^0 \left(P_{v,\text{bl}}^{(0)} - 1 \right) dt.$$

Using the symmetry (80) for $P_{v,\text{bl}}^{(0)}$ and the definition of C , we get

$$C = v_{\text{bl}}^{(2)}(0) + 2 \int_0^{+\infty} P_{v,\text{bl}}^{(0)}(t) dt, \quad (82)$$

which can be easily evaluated numerically to give $C \approx 0.8120$.

The quasipotential for $x > x_{\text{cr}}$: taking into account the rescaling of time, the equations of motion and the fact that the instanton path satisfies $H = 0$, one finds after some algebra that the quasipotential (see equation (11)) can be written as

$$\Phi_\tau = \int_{-\infty}^{+\infty} \left(\tilde{P}_v(t) \right)^2 dt, \quad (83)$$

where we denoted by $\tilde{P}_v(t)$ the instanton solution. The strategy goes as follows: we separate the contributions from the bulk and from the boundary layer in integral (83), and we evaluate both terms separately. In order to obtain a finite contribution for the boundary layer integral, we have to subtract the bulk solution from the boundary layer solution. This means that we first split the instanton path into two parts as

$$\tilde{P}_v = P_v(t) + \left(\tilde{P}_v(t) - P_v(t) \right),$$

where P_v is the bulk solution, found in section 4.1.1. We then write Φ_τ as

$$\begin{aligned} \Phi_\tau &= \int_{-\infty}^{+\infty} \left(P_v(t) + \left(\tilde{P}_v(t) - P_v(t) \right) \right)^2 dt \\ &= \int_{-\infty}^{+\infty} (P_v(t))^2 dt + 2 \int_{-\infty}^{+\infty} P_v(t) \left(\tilde{P}_v(t) - P_v(t) \right) dt + \int_{-\infty}^{+\infty} \left(\tilde{P}_v(t) - P_v(t) \right)^2 dt. \end{aligned} \quad (84)$$

Next we define $f_{\text{bl}} = \tilde{P}_v - P_v$. By construction $f_{\text{bl}} \rightarrow 0$ outside the boundary layer and f_{bl} has the same scaling as the boundary layer solution $P_{v,\text{bl}}$. More precisely f_{bl} is a natural function of $\tau'^{2/3}t$. We now turn to compute the three different contributions in the right hand side of equation (84):

Using equation (67), the bulk contribution is

$$\int_{-\infty}^{+\infty} (P_v(t))^2 dt = \int_{-\infty}^0 \left(A_0 + \frac{A_2}{\tau^{2/3}} \right)^2 e^{2t} dt = \frac{A_0^2}{2} + \frac{A_0 A_2}{\tau^{2/3}} + O\left(\frac{1}{\tau}\right) \quad (85)$$

The bulk-boundary layer cross contribution can be equivalently written using the definition of f_{bl}

$$2 \int_{-\infty}^{+\infty} P_v(t) \left(\tilde{P}_v(t) - P_v(t) \right) dt = 2 \int_{-\infty}^{+\infty} P_v(t) f(\tau'^{2/3}t) dt = \frac{2}{\tau^{2/3}} \int_{-\infty}^{+\infty} P_v(t/\tau'^{2/3}) f_{\text{bl}}(t) dt$$

Using again equation (67) and the expansion of P_v and f_{bl} to *zeroth* order we get

$$2 \int_{-\infty}^{+\infty} P_v(t) \left(\tilde{P}_v(t) - P_v(t) \right) dt \underset{\tau \rightarrow +\infty}{\sim} \frac{2A_0}{\tau^{2/3}} \int_{-\infty}^0 f_{\text{bl}}^{(0)}(t) dt + O(1/\tau),$$

where

$$f_{\text{bl}}^{(0)} = \beta \left(P_{v,\text{bl}}^{(0)}(t) - 1_{[-\infty,0]} \right).$$

Using then the symmetry of $P_{v,\text{bl}}^{(0)}(t)$ (see equation (80)), we get

$$2 \int_{-\infty}^{+\infty} P_{v,\text{B}}(t) \left(\tilde{P}_v(t) - P_{v,\text{B}}(t) \right) dt \underset{\tau \rightarrow +\infty}{\sim} - \frac{2\beta A_0}{\tau^{2/3}} \int_0^{+\infty} P_{v,\text{bl}}^{(0)}(t) dt + O(1/\tau). \quad (86)$$

The pure boundary layer contribution in equation (84) can be written

$$\begin{aligned} \int_{-\infty}^{+\infty} \left(\tilde{P}_v(t) - P_v(t) \right)^2 dt &= \int_{-\infty}^{+\infty} \left(f_{\text{bl}}(\tau^{2/3}t) \right)^2 dt \\ &\underset{\tau \rightarrow +\infty}{\sim} \frac{1}{\tau^{2/3}} \int_{-\infty}^{+\infty} \left(f_{\text{bl}}^{(0)}(t) \right)^2 dt + O(1/\tau) \\ &\underset{\tau \rightarrow +\infty}{\sim} \frac{2}{\tau^{2/3}} \int_0^{+\infty} \left(P_{v,\text{bl}}^{(0)}(t) \right)^2 dt + O(1/\tau), \end{aligned} \quad (87)$$

where we have again used in the last equality the symmetry of $P_{v,\text{bl}}^{(0)}(t)$ (see equation (80)).

Finally, summing the three contributions (85)–(87) and using the expressions for A_0 and A_2 in equations (81) and (82), we arrive to our main result

$$\forall x > x_{\text{cr}}, \quad \Phi_\tau(x) = \frac{\beta^2}{2} + \frac{\beta^2}{(\sqrt{\gamma}\beta\tau)^{2/3}} \left[v_{\text{bl}}^{(2)}(0) + 2 \int_0^{+\infty} \left(P_{v,\text{bl}}^{(0)}(t) \right)^2 dt \right] + O(1/\tau). \quad (88)$$

Moreover, the computation has shown that the bulk solution satisfies $P_v(t) = 0$ to any order in τ , for $t > 0$. This implies that the part of the instanton path after x_{cr} is a relaxation path, and has no contribution to the action (83). In particular, equation (88) is valid for any $x_{\text{cr}} < x \leq x_{\text{s}}$ outside a boundary layer of typical size $\frac{1}{\tau^{1/3}} \sqrt{\frac{\beta}{\gamma}}$ around x_{cr} .

Comments on the solution:

- (a) Both the leading order and the first correction in the limit $\tau \rightarrow \infty$ are *non-local*, that is, they only depend on the structure of $U(x)$ close to its inflection point x_{cr} , and not on the part $x > x_{\text{cr}}$. Note that if $U'(x_{\text{cr}}) = 0$, using similar arguments as was done to find the scaling of the boundary layer, one expects a correction scaling as $\frac{1}{\tau^{4/7}}$ instead of $\frac{1}{\tau^{2/3}}$.
- (b) The correction at order $\frac{1}{\tau^{2/3}}$ given by equation (88) has two origins. The first contribution comes from the fact that, for finite τ , the Ornstein–Uhlenbeck process has to overshoot the velocity fluctuation above β . The fluctuation of the Ornstein–Uhlenbeck process up to the value $\beta + \frac{\beta}{\tau^{2/3}} v_{\text{bl}}^{(2)}(0)$ is simply $\frac{\left(\beta + \frac{\beta}{\tau^{2/3}} v_{\text{bl}}^{(2)}(0) \right)^2}{2}$

$= \frac{\beta^2}{2} + \frac{\beta^2}{\tau^{2/3}} v_{\text{bl}}^{(2)}(0) + O(1/\tau^{4/3})$. The integral correction in equation (88) can be interpreted as the price to pay for sustaining the active velocity at the value β during the crossing of the inflection point. It thus comes from the fact that crossing the inflection point takes a finite time, when τ is finite.

- (c) Whereas the correction at order $\frac{1}{\tau^{1/3}}$ vanishes, there is no reason for the correction at order $\frac{1}{\tau}$ to be zero, because the bulk instanton part has a contribution to this order. One can thus expect the next order to scale as $\frac{1}{\tau}$.
- (d) A numerical resolution of the boundary layer equations (74) and (75) gives $v_{\text{bl}}^{(2)}(0) \approx 0.5905$ and $2 \int_0^{+\infty} \left(P_{v,\text{bl}}^{(0)}(t) \right)^2 dt = 0.2215$. The sum of these two values gives the numerical constant C of the result announced in equation (4). For the escape problem ($x = x_{\text{cr}}$), our result coincide with that of [7], including for the numerical estimate of the constant C (note also that the dominant order of (4) was derived earlier in [2]). In [7] the perturbation theory presents no boundary layer.

4.2. The quasipotential for $x < x_{\text{cr}}$

We now turn to the computation of the quasipotential for $x < x_{\text{cr}}$. In the following, we only sketch the computations as they are very similar to the case $x > x_{\text{cr}}$. We look for an instanton path solving equations (56) and (57) with the boundary conditions equation (14).

4.2.1. Bulk solution. The argument developed in section 4.1.1 to show that $P_x = 0$ to any order in τ remains valid in the range $x < x_{\text{cr}}$. The instanton equations in the bulk are

$$\begin{cases} \dot{x} = \tau (v - U'(x)), \\ \dot{v} = -v + 2P_v, \\ \dot{P}_v = P_v. \end{cases} \quad (89)$$

Contrary to the domain $x > x_{\text{cr}}$, there is no anomalous scaling for the boundary layer in the part $x < x_{\text{cr}}$. We thus expand the solution as

$$P_v(t) = P_v^{(0)}(t) + \frac{1}{\tau} P_v^{(1)}(t) + \frac{1}{\tau^2} P_v^{(2)}(t) + \dots, \quad (90)$$

and similarly for $x(t)$ and $v(t)$. Thanks to the linearity of equation (89) for P_v and v , the solution to any order k can be written

$$t < 0: \begin{cases} v^{(k)}(t) = A_k e^t, \\ P_v^{(k)}(t) = A_k e^t. \end{cases} \quad (91)$$

One can then find the solution $x^{(k)}(t)$ using an expansion of $\dot{x} = \tau (v - U'(x))$ to k th order in $\frac{1}{\tau}$. The constants A_k have to be determined by matching the bulk solution to the boundary layer as we do below to up to first order.

4.2.2. Boundary-layer solution. The boundary layer equations are obtained from equations (56) and (57) by rescaling time according to $t' = \tau t$ and expanding U' in the vicinity of x as

$$U'(x') = \beta + \gamma(x' - x), \quad (92)$$

where $\beta = U'(x)$ and $\gamma = U''(x)$. Note that both are strictly positive. We then define the new variable $y = x' - x$ and we get the boundary layer equations

$$\begin{cases} \dot{y}_{\text{bl}} = v_{\text{bl}} - \gamma y_{\text{bl}}, \\ \dot{v}_{\text{bl}} = -\frac{v_{\text{bl}}}{\tau} + 2\frac{P_{v,\text{bl}}}{\tau}, \end{cases} \quad (93)$$

and

$$\begin{cases} \dot{P}_{x,\text{bl}} = \gamma P_{x,\text{bl}}, \\ \dot{P}_{v,\text{bl}} = \frac{P_{v,\text{bl}}}{\tau} - P_{x,\text{bl}}. \end{cases} \quad (94)$$

We then expand the boundary layer solution as

$$y_{\text{bl}}(t) = y_{\text{bl}}^{(0)}(t) + \frac{1}{\tau} y_{\text{bl}}^{(1)}(t) + \frac{1}{\tau^2} y_{\text{bl}}^{(2)}(t) + \dots$$

and use similar expansions for $P_{x,\text{bl}}, P_{v,\text{bl}}, v_{\text{bl}}(t)$. We now turn to solve these equations order by order.

Zeroth order: to this order we get

$$\begin{cases} y_{\text{bl}}^{(0)}(t) = 0 \\ v_{\text{bl}}^{(0)}(t) = \beta \\ P_{v,\text{bl}}^{(0)} = \beta(1 - e^{\gamma t}). \end{cases} \quad (95)$$

Matching this with the bulk solution of equation (91) implies $A_0 = \beta$.

First order: to order $1/\tau$ we get

$$\begin{cases} v_{\text{bl}}^{(1)}(t) = \beta t + \frac{2\beta}{\gamma}(1 - e^{\gamma t}), \\ P_{v,\text{bl}}^{(1)} = \beta t + \frac{2\beta}{\gamma}(1 - e^{\gamma t}), \end{cases} \quad (96)$$

and matching to the bulk solution of equation (91) implies $A_1 = \frac{2\beta}{\gamma}$. Note that we do not need to derive an explicit expression for $y_{\text{bl}}^{(1)}$ in order to calculate the quasipotential.

The quasipotential for $x < x_{\text{cr}}$: using the same decomposition as in equation (83) and in the discussion that follows, we can express the quasipotential to order $1/\tau$ as

$$\begin{aligned} \Phi_\tau(x) &\underset{\tau \rightarrow +\infty}{\sim} \int_{-\infty}^0 (P_v^{(0)}(t))^2 dt + \frac{2}{\tau} \int_{-\infty}^0 P_v^{(0)}(t) P_v^{(1)}(t) dt \\ &+ \frac{2}{\tau} \int_{-\infty}^0 P_v^{(0)}(t) f_{\text{bl}}^{(0)}(t) dt + \frac{1}{\tau} \int_{-\infty}^0 (f_{\text{bl}}^{(0)}(t))^2 dt, \end{aligned} \quad (97)$$

where we have used again the notation $f_{\text{bl}} = \tilde{P}_v - P_v$, with \tilde{P}_v the instanton path and P_v the bulk solution. A straightforward evaluation of the different terms in equation (97) using the solutions equations (91), (95) and (96) leads to the final result announced in equation (3), namely

$$\Phi_\tau(x) \underset{\tau \rightarrow +\infty}{\sim} \frac{\beta^2}{2} + \frac{\beta^2}{2\gamma\tau} + O(1/\tau^2). \quad (98)$$

5. Perspectives

We derived the escape rate and the stationary distribution of an AOUP, in expansions at small and large correlation time τ . By focusing on the asymptotics of small activity (in amplitude), we were able to use exact large-deviation techniques (i.e. without using the UCNA or the Fox approximation schemes that are often employed to study systems with colored noise). In the $\tau \rightarrow 0$ limit the quasipotential is local to first order, as already known, but becomes non-local at second order in τ . In the large τ limit the quasipotential is inherently non-local and singular. This leads to a host of physical consequences such as a ratchet effect and a possible fly-over of metastable state facilitated by the memory of the active noise.

The results we have described open many questions. The path-integral technique used enables one to derive the quasipotential as a functional of an optimal trajectory of an equilibrium problem; as we focused, for simplicity, on the one-dimensional case, the explicit dependence on the time-dependent trajectory could be eliminated [see for instance the passage from (21) to (22)]. This implies that the quasipotential can be reexpressed as a functional of the potential only and not of the optimal trajectory. The generalization of our computation to higher dimensions is rather immediate, but it is not obvious that at the last stage one can similarly eliminate the explicit dependence on the optimal trajectory. This could potentially lead to interesting effects in cases where several optimal trajectories are in competition.

We have seen that the small- τ and the large- τ asymptotics present very different physical features: at small τ , the optimal trajectory remains close to the equilibrium one (that bypasses a potential barrier), while at large τ the dynamics is dominated by a ‘barrier of force’ (i.e. the region of maximal force). The switch from a potential to a force barrier dominated regime could either be a cross-over or signal a singularity as τ is increased. However, our perturbative approach leaves this question open. Last, since our technique allows for a treatment of generic colored-noise dynamics (in the small noise limit), it could be instructive to compare its predictions to those of the UCNA and the Fox approximations in a systematic way.

Acknowledgments

We thank Nir Gov for his involvement in a project that led to this work, and an anonymous referee for bringing our attention to references [4, 6, 7]. E W and Y K are supported by and ISF Grant and an NSF-BSF Grant. E W is partly supported by a

Technion Grant. V L thanks the Technion, where part of this work was undergone, for kind hospitality. V L is supported by the ERC Starting Grant No. 680275 MALIG, the ANR-18-CE30-0028-01 Grant LABS and the ANR-15-CE40-0020-03 Grant LSD.

Appendix. Mean escape time from a metastable state

Let us now recall how the quasipotential is related to the mean escape time from a metastable state x_0 . Again in the following, x_s is the saddle point to escape from the attraction basin of x_0 . We show how the mean escape time $\langle T_{\text{esc}} \rangle$ can be computed directly from the quasipotential in the limit where D is small.

The standard procedure is to compute the mean escape time $\langle T_{\text{esc}} \rangle$ from the Fokker–Planck equation with absorbing boundary conditions: let $J(t)$ be the outgoing flux at $x = x_s$, we have simply

$$\langle T_{\text{esc}} \rangle = \int_0^{+\infty} t J(t) dt. \quad (99)$$

in general, there is no simple solution to this problem. But one can derive the result when escape is a rare event, which correspond to the small D limit in equation (1). In this limit, the outgoing flux is exponentially small in $1/D$, such that the relaxation time T_{rel} of the particle inside the trap is much smaller than the mean escape time $\langle T_{\text{esc}} \rangle$. A good approximation is then to consider that two consecutive escapes are independent events, which means that escape is a Poisson process with rate λ . On the timescale $t \gg T_{\text{rel}}$, the probability $M(t)$ that the particle is still in the trap within the interval $[0, t]$ follows the equation

$$\frac{dM}{dt} = -\lambda M.$$

This relation can be immediately integrated to give

$$M(t) = e^{-\lambda t}.$$

$J(t)$ is related to $M(t)$ through the simple relation

$$J(t) = -\frac{dM}{dt} = \lambda e^{-\lambda t}. \quad (100)$$

It follows from equations (99) and (100) that the mean escape time satisfies

$$\frac{1}{\langle T_{\text{esc}} \rangle} = \lambda = \lim_{T_{\text{rel}} \ll t \ll \langle T_{\text{esc}} \rangle} J(t). \quad (101)$$

On the other hand, J is related to the transition probability through

$$J(t) = P(T_{\text{esc}} = t) = P(x_s, t | x_0, 0). \quad (102)$$

In the intermediate timescale regime $T_{\text{rel}} \ll t \ll \langle T_{\text{esc}} \rangle$, the transition probability $P(x_s, t|x_0, 0)$ does not depend on t and is given by the large deviation principle

$$P(x_s, t|x_0, 0) \underset{D \rightarrow 0}{\asymp} e^{-\frac{\Phi_T(x_s)}{D}}. \quad (103)$$

Relations (101)–(103) together imply the large deviation result

$$\langle T_{\text{esc}} \rangle \underset{D \rightarrow 0}{\asymp} e^{\frac{\Phi_T(x_s)}{D}}. \quad (104)$$

References

- [1] Pesquera L, Rodriguez M A and Santos E 1983 Path integrals for non-Markovian processes *Phys. Lett. A* **94** 287
- [2] Luciani J F and Verga A D 1987 Functional integral approach to bistability in the presence of correlated noise *Europhys. Lett.* **4** 255
- [3] Klocek-Dygas M M, Matkowsky B J and Schuss Z 1988 Colored noise in dynamical systems *SIAM J. Appl. Math.* **48** 425
- [4] Bray A J and McKane A J 1989 Instanton calculation of the escape rate for activation over a potential barrier driven by colored noise *Phys. Rev. Lett.* **62** 493
- [5] Peter H 1989 Path integral solutions for non-Markovian processes *Z. Phys. B* **75** 275
- [6] McKane A J, Luckock H C and Bray A J 1990 Path integrals and non-Markov processes: I. General formalism *Phys. Rev. A* **41** 644
- [7] Bray A J, McKane A J and Newman T J 1990 Path integrals and non-Markov processes: II. Escape rates and stationary distributions in the weak-noise limit *Phys. Rev. A* **41** 657
- [8] Luckock H C and McKane A J 1990 Path integrals and non-Markov processes: III. Calculation of the escape-rate prefactor in the weak-noise limit *Phys. Rev. A* **42** 1982
- [9] Jung P and Peter H 1987 Dynamical systems: a unified colored-noise approximation *Phys. Rev. A* **35** 4464
- [10] Hänggi P and Jung P 1995 Colored noise in dynamical systems *Adv. Chem. Phys.* **89** 239
- [11] Szamel G 2014 Self-propelled particle in an external potential: existence of an effective temperature *Phys. Rev. E* **90** 012111
- [12] Koumakis N, Maggi C and Di Leonardo R 2014 Directed transport of active particles over asymmetric energy barriers *Soft Matter* **10** 5695
- [13] Farage T F F, Krinninger P and Brader J M 2015 Effective interactions in active Brownian suspensions *Phys. Rev. E* **91** 042310
- [14] Szamel G, Flenner E and Berthier L 2015 Glassy dynamics of athermal self-propelled particles: computer simulations and a nonequilibrium microscopic theory *Phys. Rev. E* **91** 062304
- [15] Fodor É, Nardini C, Cates M E, Tailleur J, Visco P and van Wijland F 2016 How far from equilibrium is active matter? *Phys. Rev. Lett.* **117** 038103
- [16] Caprini L, Marconi U M B, Puglisi A and Angelo V 2019 Active escape dynamics: the effect of persistence on barrier crossing *J. Chem. Phys.* **150** 024902
- [17] Woillez E, Kafri Y and Gov N 2020 Active trap model *Phys. Rev. Lett.* **124** 118002
- [18] Lambert G, Liao D and Austin R H 2010 Collective escape of chemotactic swimmers through microscopic ratchets *Phys. Rev. Lett.* **104** 168102
- [19] Di Leonardo R, Angelani L, Dell Arciprete D, Ruocco G, Iebba V, Schippa S, Conte M P, Mecerini F, De Angelis F and Di Fabrizio E 2010 Bacterial ratchet motors *Proc. Natl Acad. Sci.* **107** 9541
- [20] Angelani L, Costanzo A and Di Leonardo R 2011 Active ratchets. *Europhys. Lett.* **96** 68002
- [21] Wio H S, Colet P, San Miguel M, Pesquera L and Rodríguez M A 1989 Path-integral formulation for stochastic processes driven by colored noise *Phys. Rev. A* **40** 7312
- [22] Newman T J, Bray A J and McKane A J 1990 Inertial effects on the escape rate of a particle driven by colored noise: an instanton approach *J. Stat. Phys.* **59** 357
- [23] McKane A J 1989 Noise-induced escape rate over a potential barrier: results for general noise *Phys. Rev. A* **40** 4050
- [24] Freidlin M I and Wentzell A D 2012 *Random Perturbations of Dynamical Systems* 3rd edn (New York: Springer)

- [25] Baek Y, Solon A P, Xu X, Nikola N and Kafri Y 2018 Generic long-range interactions between passive bodies in an active fluid *Phys. Rev. Lett.* **120** 058002
- [26] Grafke T and Vanden-Eijnden E 2017 Non-equilibrium transitions in multiscale systems with a bifurcating slow manifold *J. Stat. Mech.* **093208**
- [27] Onsager L and Machlup S 1953 Fluctuations and irreversible processes *Phys. Rev.* **91** 1505
- [28] Cugliandolo L F, Lecomte V and van Wijland F 2019 Building a path-integral calculus: a covariant discretization approach *J. Phys. A: Math. Theor.* **52** 50LT01
- [29] Janssen H-K 1976 On a Lagrangean for classical field dynamics and renormalization group calculations of dynamical critical properties *Z. Phys. B* **23** 377
- [30] Janssen H-K 1979 Field theoretical methods applied to critical dynamics *Dynamical Critical Phenomena and Related Topics* (Lecture Notes in Physics vol 104) ed C P Enz (Berlin: Springer) p 26
- [31] De Dominicis C 1976 Techniques de renormalisation de la théorie des champs et dynamique des phénomènes critiques *J. Phys. Colloq.* **37** C1
- [32] De Dominicis C and Peliti L 1978 Field-theory renormalization and critical dynamics above T_c : helium, antiferromagnets, and liquid-gas systems *Phys. Rev. B* **18** 353
- [33] Martin P C, Siggia E D and Rose H A 1973 Statistical dynamics of classical systems *Phys. Rev. A* **8** 423
- [34] Baek Y and Kafri Y 2015 Singularities in large deviation functions *J. Phys. A: Math. Theor.* **2015** P08026